Georgian Scientists
ქართველი მეცნიერები
Vol. 8 Issue 4(1), 2026
https://doi.org/10.52340/gs.2026.08.04(1).115



PELVIC FRACTURES

Vazha Bardavelidze

Caucasus Medical Centre, Tbilisi, Georgia

Background and context: Pelvic fractures represent some of the most severe and complex injuries encountered in emergency medicine. Modern treatment of these fractures requires a highly coordinated, multi-disciplinary approach designed to minimize initial mortality and optimize long-term physiological recovery.

Problem statement: The acute management phase poses critical challenges, primarily due to the high risk of catastrophic hemorrhage, severe hemodynamic instability, and associated intra-abdominal or retroperitoneal injuries. Managing these life-threatening conditions during the golden hour remains a paramount hurdle in emergency trauma care.

Methodology and approach: To treat these high-energy injuries properly, a standardized, aggressive protocol must be initiated immediately upon admission. This approach integrates advanced damage control resuscitation with rapid mechanical stabilization using pelvic binders or temporary external fixation. When indicated, immediate surgical interventions such as pre-peritoneal pelvic packing or angioembolization are deployed to achieve definitive hemorrhage control, followed by timed anatomical reconstruction.

Key results: Utilizing this structured acute management framework drastically minimizes early mortality rates. Furthermore, early rigid stabilization enables accelerated patient mobilization and early rehabilitation. This proactive timeline actively prevents debilitating post-traumatic complications—such as deep vein thrombosis, prolonged pulmonary dysfunction, and chronic pelvic instability—leading to the best possible functional outcomes.

Conclusion: The optimal treatment strategy for acute pelvic trauma relies heavily on swift, multidisciplinary damage control protocols. Combining immediate mechanical and physiological stabilization with early, aggressive rehabilitation forms the definitive gold standard for maximizing patient survival and restoring baseline quality of life.

Keywords: Pelvic fractures, acute trauma, hemorrhage control, early rehabilitation, damage control.